



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From which immediately $x=2$, $y=-2$.

Dividing we have $x+2 = -\frac{1}{2-y}$, or $y = \frac{2x+5}{x+2}$.

Substitute in (1), and obtain $x^3+2x^2+1=0$. Solve this by any of the above methods.

V. Add (1) and (2) and complete squares. Then

$$x^2+x+\frac{1}{4}+y^2+y+\frac{1}{4}=8+\frac{2}{4} \text{ or } (x+\frac{1}{2})^2+(y+\frac{1}{2})^2=\frac{25}{4}+\frac{9}{4}.$$

Whence, from the four possible corresponding values of x and y , we may pick out one set which satisfies the original system, namely $x=2$, $y=-2$.

Also solved by *GEO. R. BERRY*.

NOTE. The donor of this prize has acted as judge on the merits of the several solutions, and his decision is that the two published solutions are of equal merit. In accordance with this decision, the prize money has been equally divided between Miss Scheffer and Mr. Young. We might say that there has been only one solution sent in to the prize problem in Mechanics. This solution is defective. The problem is, therefore, open to all our contributors for solution. **EDITOR F.**

105. Proposed by *CHARLES E. MYERS*, Canton, O.

Solve for x the following: $a \log(ax^2) = m \log(m)$.

Solution by *G. B. M. ZERR*, A. M., Ph. D., Professor of Science and Mathematics, Chester High School, Chester, Pa.; *J. K. ELLWOOD*, A. M., Colfax School, Pittsburg, Pa.; *J. SCHEFFER*, A. M., Hagerstown, Md.; *COOPER D. SCHMITT*, A. M., University of Tennessee, Knoxville, Tenn.; *W. F. SHAW*, Austin, Tex.; and *ELMER SCHUYLER*, B. Sc., Professor of German and Mathematics, Boys' High School, Reading, Pa.

$a \log(ax^2) = m \log m$ may be written $a^{x^2 \log a} = m^{\log m}$.

$$\therefore x^2 (\log a)^2 = (\log m)^2.$$

$$\therefore x = \pm (\log m / \log a).$$

GEOMETRY.

130. Proposed by *B. F. FINKEL*, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.

If the points x , y , z divide the strokes $c-b$, $a-c$, $b-a$, in the same ratio r , and the triangles x , y , z and a , b , c are similar, either $r=1$ or both triangles are equilateral. [From Harkness and Morley's *Introduction to the Theory of Functions*, page 26].

Solution by the **PROPOSER**.

Let x , y and z denote the points, dividing $c-b$, $a-c$, and $b-a$, respectively, in the given ratio r .

$$\text{Then } x = \frac{c+br}{1+r}, \quad y = \frac{a+cr}{1+r}, \quad \text{and } z = \frac{b+ar}{1+r}.$$

The condition that a , b , c , and x , y , z form similar triangles is

$$\begin{vmatrix} a & x & 1 \\ b & y & 1 \\ c & z & 1 \end{vmatrix} = 0 \dots (1).$$